



## **EURAP**

# **An International Antiepileptic Drugs and Pregnancy Registry**

Interim Report – MAY 2021

### **Central Study Coordinator**

Dr. Dina Battino

*Department of Neurophysiology and Experimental Epileptology*

*Fondazione IRCCS Istituto Neurologico Carlo Besta*

*20 133 Milano, Italy*

Tel: + 39-02-23-94-22-30

Fax: + 39-02-700-42-91-60

E-mail: [eurap@istituto-besta.it](mailto:eurap@istituto-besta.it)

### **Chairman Central Project Commission**

Prof. Torbjörn Tomson

*Department of Neurology*

*Karolinska University Hospital*

*S-171 76 Stockholm, Sweden*

Tel: + 46-08-51-77-37-05

Fax: + 46-08-51-77-37-57

E-mail: [torbjorn.tomson@sll.se](mailto:torbjorn.tomson@sll.se)

## **BACKGROUND**

A number of independent groups with experience and interest in maternal and foetal well-being in association with maternal use of antiepileptic drugs (AEDs) have agreed on a prospective international multi-centre study of pregnancies with AEDs. Data from all participating groups are shared in a Central Registry of Antiepileptic Drugs and Pregnancy (EURAP). EURAP was established in the first centres in some European countries and has since then gradually expanded to include more centres and countries now involving also Asia, Oceania, Latin America and Africa.

## **OBJECTIVE OF EURAP**

The primary objective of EURAP is to evaluate and determine the comparative risk of major foetal malformations following intake of AEDs and their combinations during pregnancy.

## **METHODS**

EURAP is an observational study. Women taking AEDs at the time of conception, irrespective of the indication, may be included. To avoid selection bias, only pregnancies recorded before foetal outcome is known and within week 16 of gestation are included in the prospective risk assessment. Cases ascertained later in pregnancy are recorded as retrospective cases, as they may provide signals, but are not included in the comparative risk evaluation.

Information on patient's demographics, type of epilepsy, seizure frequency, family history of malformations, drug therapy and of other potential risk factors is obtained, and follow-up data are collected once at each trimester, at birth and at one year after delivery.

Networks of reporting physicians have been established in countries taking part in the collaboration. During the course of the pregnancy, and the follow-up time after delivery, the participating physician enters data into five Subforms (Subforms A-E) for each patient.

Subform A is completed on enrolment of the patient, Subform B after the first trimester, Subform C after the second trimester, Subform D within three months after delivery, and Subform E within 14 months after birth. Immediately after completion, each Subform is submitted to the national coordinator for review. The national coordinator transfers the reviewed and accepted Subform to the Central EURAP Registry in Milan, Italy.


## EVALUATION OF OUTCOME

The physician records descriptively abnormalities observed in the offspring. The final assessment and classification of the type of malformation is the responsibility of the Central Project Commission (CPC). In order to facilitate a uniform and objective assessment, reports of malformations are assessed regularly by an outcome assessment committee, which is kept blinded with respect to the type of exposure.

## INTERIM REPORT

EURAP was implemented in the first two countries in Europe in 1999 and has since then grown to include countries from Europe, Oceania, Asia, Latin America and Africa. This development is reflected by increasing numbers of enrolled pregnancies. The development since 1999 is illustrated in Fig. 1.

**Fig 1. Number of Participating Countries and Pregnancies Reported to the Central Registry by March, 2021.**



The present report is based on data available in the Central Registry by May 11th, 2021.

At that time more than 1,500 reporting physicians from 45 countries had contributed cases to the Central Registry. Countries that have contributed at least 10 pregnancies in the current report are listed in Table 1.

**Table 1. Countries that have contributed at least 10 pregnancies in the current report (n=36).**

<b>COUNTRY</b>	<b>National Coordinator (or referring physician*)</b>	<b>Date of joining the Registry</b>
Argentina	Silvia Kochen	2002
Australia	Frank Vajda	2000
Austria	Gerhard Luef	2000
Belgium	Dick Lindhout & Eugène van Puijenbroek	2002
Chile	Alejandro De Marinis	2002
China	Weiping Liao	2006
Croatia	Dinko Vitezic	2002
Czech Republic	Jana Zarubova	2001
Denmark	Anne Sabers	2000
El Salvador	Ovidio Solano Cabrera*	2017
Finland	Reetta Kälviäinen	2003
France	Aileen McGonigal*	2000
Georgia	Sofia Kasradze; Nino Gogarishvili*	2000
Germany	Bettina Schmitz	2000
Hong-kong	Patrick Kwan	2002
India	Sanjeev Thomas	2001
Iran	Nasim Tabrizi	2018
Israel	Lilach Goldstein	2000
Italy	Luigi M. Specchio	2000
Japan	Hideyuki Ohtani	2001
Lithuania	Ruta Mameniskiene	2002
Macedonia	Gordana Kiteva Trancevska	2001
Netherlands	Dick Lindhout & Eugène van Puijenbroek	2002
Norway	Silje Alvestad	2000
Philippines	Leonor Cabral-Lim	2003
Poland	Joanna Jedrzejszak	2001
Portugal	Isabel Pires*; Joana Parra*; Ines Cunha*; Elia Baeta*; Carla Bentes*; Catarina Cruto*; Inês Menezes Cordeiro*	2001
Serbia & Montenegro	Maja Milovanovic	2002
Slovakia	Vladimír Safcák	2002
Slovenia	Boštjan Čebular & Gal Granda	2002
Spain	Meritxell Martinez Ferri	2001
Sweden	Torbjörn Tomson	2000
Switzerland	Barbara Tettenborn & Elisabeth Langenberger; Dominique Flügel*	2001
Taiwan	Hsiang-Yu Yu	2004
Turkey	Demet Ilhan Algin	2000
United Kingdom	John Craig & Craig Heath	2001

\* referring physicians

By the cut-off date for this report (May, 11th, 2021), **27,896 pregnancies had been entered into the central database**. Of these, **12,135 pregnancies are excluded** from the present interim report for reasons explained here below:

1. Pregnancies that failed to meet inclusion criteria (n=201).
2. Lost to follow-up, including those failing to submit sub-forms within preset deadlines (n=3,628).
3. Pending pregnancies, awaiting updates or corrections of different sub-forms (n=1,308).
4. Ongoing pregnancies, updated and corrected (n=630).
5. Retrospective, but completed and corrected (n=4,541). Among these, there are true retrospective pregnancies (n=4,188) and a further three hundred and fifty three pregnancies (n=353) that otherwise met our criteria for prospective pregnancies since they were recruited within 16<sup>th</sup> week, but for which patients had an ultrasound examination performed before enrolment.
6. Retrospective, i.e. initially classified as prospective pregnancies but re-classified as retrospective cases because one or more CRF subforms were submitted after the set deadlines (n=402).
7. Unclassifiable i.e. cases for which it was impossible to determine if there was a malformation or not (n=90). This includes 1 stillbirth with unknown fetal status, induced abortions with insufficient information on fetus (n=6), anomalies in livebirths where the information was insufficient to determine if qualifying for malformation diagnosis (n=78), 1 incomplete spontaneous abortion with unclear results of biopsy, and 4 perinatal deaths in premature births (<35 gestational weeks) with anomalies difficult to classify as congenital or due to prematurity.
8. Not yet classified, i.e. pregnancies which classification is pending as well as pregnancies which became completed after the last time we sent the database to the Outcome Assessment Committee (OAC), regardless if they contained some malformations or not (n=216).
9. Treatment changes between different AEDs or mono- to polytherapy or vice versa during the first trimester (n=1,119).

Thus in total **15,761 prospective pregnancies** (enrolled at the latest during the 16<sup>th</sup> gestational week and before outcome is known) **are included** in this report.

The classification of the epilepsy among the prospective pregnancies is given in table 2. Epilepsy was the indication for treatment in all but 118 (0.8%) of the pregnant women.

**Table 2. Classification of the epilepsy in 15,761 prospective pregnancies.**

<b>Epilepsy</b>	<b>N</b>	<b>%</b>
Localisation-related*	8,258	52.4
Generalized	6,563	41.6
Undetermined	517	3.3
Missing information	305	1.9
No epilepsy	118	0.8
<b>Total</b>	<b>15,761</b>	<b>100</b>

*\*Focal, according to more current terminology.*

The **maternal age** among prospective cases was **30.1 ±5.1 years** (mean±SD), ranging from 14 to 55 years.

The women were of Caucasian **ethnicity** in 87% and of Asian in 10%.


**Gravida for each pregnancy** is presented in Table 3.

**Table 3. Number of the pregnancy in prospective cases.**

<b>Gravida</b>	<b>N</b>	<b>%</b>
1st pregnancy	7,156	45.5
2nd pregnancy	4,939	31.3
3rd pregnancy	2,180	13.8
4th pregnancy	905	5.7
5th pregnancy	357	2.3
> 5th pregnancy	221	1.4
Not ascertained	3	0.0
<b>Total</b>	<b>15,761</b>	<b>100</b>


The outcome of the prospective completed pregnancies is presented in Figure 2. Out of the **294 induced abortions**, 47 were for chromosomal abnormalities and/or syndromes and 77 were for other fetal indication detected by prenatal screening (*out of these 77 cases, 62 were confirmed as major malformations and the remaining 15 cases were definitively classified as other abnormalities such as hydrops fetalis, molar pregnancies, blighted ovum, fetal placental transfusion syndromes, fetal growth retardation, fetus papyraceus, fetal death for unspecified causes, balanced translocation and insertion in normal individual, Dravet syndrome*).

Figure 2. Obstetrical outcome of prospective pregnancies.



Of the pregnancies, **12,579 (79.8%)** involved women on a single AED, 2,560 (16.2%) were on two AEDs whereas 448 (2.9%) took three AEDs or more. One hundred and seventy-four women (1.1%) were not on AED treatment during the 1<sup>st</sup> trimester. The exposure to the different AEDs in monotherapy among the prospective pregnancies is presented in Figure 3.

Figure 3. Number of prospective pregnancies with exposure to different AEDs in monotherapy during the first trimester of pregnancy.



**There were 334 different AED combinations.** The most frequently used combinations were lamotrigine and levetiracetam (n=371), lamotrigine and valproic acid (n=287), carbamazepine and levetiracetam (n=170), carbamazepine and clobazam (n=128), carbamazepine and lamotrigine (n=125), lamotrigine and topiramate (n=101), carbamazepine and valproic acid (n=84), carbamazepine and phenobarbital (n=83), clobazam and lamotrigine (n=63), levetiracetam and oxcarbazepine (n=63), levetiracetam and valproic acid (n=60), clonazepam and lamotrigine (n=57), and carbamazepine and topiramate (n=55) (Table 4).

**Table 4. The most common AED combinations.**

<b>The most common polytherapies during the first trimester of pregnancy</b>	<b>N</b>
lamotrigine + levetiracetam	371
lamotrigine + valproic acid	287
carbamazepine + levetiracetam	170
carbamazepine + clobazam	128
carbamazepine + lamotrigine	125
lamotrigine + topiramate	101
carbamazepine + valproic acid	84
carbamazepine + phenobarbital	83
clobazam + lamotrigine	63
levetiracetam + oxcarbazepine	63
levetiracetam + valproic acid	60
clonazepam + lamotrigine	57
carbamazepine + topiramate	55
lamotrigine + oxcarbazepine	44
clonazepam + valproic acid	40
topiramate + valproic acid	39
phenobarbital + valproic acid	38
levetiracetam + topiramate	35
phenobarbital + phenytoin	33
carbamazepine + clonazepam	33
clobazam + oxcarbazepine	32
lacosamide + levetiracetam	28
lamotrigine + phenobarbital	27

The number of pregnancies with exposure to different second generation AEDs taken in combination with other AEDs are listed in Table 5.

**Table 5. Number of pregnancies with different second generation AEDs in combination therapy.**

Lamotrigine	1,392
Levetiracetam	1,011
Topiramate	388
Oxcarbazepine	266
Zonisamide	91
Lacosamide	81
Gabapentin	65
Vigabatrin	37
Pregabalin	29
Eslicarbazepine acetate	20
Tiagabine	11
Perampanel	10
Brivaracetam	5
Rufinamide	2
Retigabine	1

## TERATOGENIC OUTCOME

There were 702 major congenital malformations (MCM), 24 syndromic and/or genetic cases and 88 chromosomal abnormalities (CHR) in the prospective cohort of 14,875 pregnancies as shown in Table 6 (886 spontaneous abortions are excluded).

**Table 6. Pathological outcomes.**

Outcome	Outcome Classification	N
MCM	Multiple major	57
	Isolated major	645
<b>MCM</b>		<b>702</b>
<b>SYNDROMES or GENETIC conditions</b>		<b>24</b>
<b>CHR</b>		<b>88</b>
<b>Total</b>		<b>814</b>

The 24 syndromic and/or genetic cases are Marfan's syndrome (3), Noonan syndrome (3), inherited tuberous sclerosis (6), Goldenhar syndrome (1), incontinentia pigmenti (2), inherited congenital glaucoma (1), inherited congenital cataract (1), inherited craniosynostosis (1), Di George's syndrome (1), bilateral hearing loss (1), X-linked lissencephaly (1), Skeletal dysplasia/Dwarfism (1), X-linked ichthyosis (1) and Freeman Sheldon syndrome (1).

In this report we will confine our analysis to the 702 MCM including 62 induced abortions, six stillbirths and 16 neonatal deaths. Of the 618 live births, 83 cases of malformations were ascertained prenatally, 366 were first reported at birth, and a further 169 cases not detected at birth but within one year after birth.

Among the 702 cases with MCM, 154 were detected by ultrasound examination. Out of these 154 cases, there were 62 induced abortions, five stillbirths, four perinatal deaths and 83 live births.

The 702 cases represent a [malformation prevalence of 4.7%](#) of all prospective pregnancies for which follow-up has been completed (702/14,875).

**The type of malformations is described in Table 7.**

**Table 7**

<b>PATHOLOGICAL OUTCOMES</b>	<b>DESCRIPTION</b>	<b>N</b>
MCM	Multiple major	57
	<b>Nervous system</b>	
MCM	Spina Bifida	42
MCM	Anencephalus and similar	4
MCM	Hydrocephaly	7
MCM	Microcephaly	2
MCM	Nervous system (other malformations)	16
	all	71
	<b>Cardiovascular system</b>	
MCM	Atrial septal defect	37
MCM	Ventricular septal defect	61
MCM	Atrioventricular septal defect	2
MCM	Congenital heart disease	54
MCM	Tetralogy of Fallot	4
MCM	Transposition of great vessels (complete)	4
MCM	Pulmonary valve stenosis	10
MCM	Hypoplastic left heart	8
	all	180
	<b>Urinary system</b>	
MCM	Urinary system (other malformations)	51
MCM	Renal Dysplasia	7
	all	58
	<b>Digestive system</b>	
MCM	Diaphragmatic hernia	8
MCM	Ano-rectal atresia and stenosis	2
MCM	Digestive system (other malformations)	11
MCM	Duodenal atresia or stenosis	3
MCM	Gastroschisis	3
MCM	Omphalocele	3
MCM	Atresia of oesophagus without fistula	2
	all	32
	<b>Limbs</b>	
MCM	Upper limb reduction	8
MCM	Lower limb reduction	1
MCM	Syndactyly	8
MCM	Polydactyly	24
MCM	Club foot - talipes equinovarus	22
	all	63
	<b>Musculoskeletal</b>	
MCM	Musculo-skeletal (other malformations)	11
MCM	Hip dislocation and/or dysplasia	70
	all	81
	<b>Genital system</b>	
MCM	Genital (developmental ovarian cyst)	6
MCM	Hypospadias	80
	all	86
	<b>Eye, ear, face and neck</b>	
MCM	Congenital cataract	4
MCM	Eye (other malformations)	3
MCM	Ear, face and neck	5
MCM	Choanal atresia	1
MCM	Atresia of nasopharynx	1
	all	14
	<b>Oro facial clefts</b>	
MCM	Cleft lip with or without palate	15
MCM	Cleft palate	16
	all	31
MCM	Other specified malformations (including sacral teratoma, cystic hygroma, haemangiomas, accessory skin tags, aberrant subclavian artery, congenital malformation of spleen, sequences, genetic syndromes, congenital malformation of renal artery, congenital malformation of adrenal gland, congenital malformations of integument, congenital malformations of lung, congenital bronchomalacia)	29
<b>MCM</b>	<b>all MCMs</b>	<b>702</b>
<b>CHR</b>	<b>all CHR</b>	<b>88</b>
<b>Syndromes</b>	<b>all Syndromes</b>	<b>24</b>
<b>Total</b>	<b>all cases with pathological outcomes</b>	<b>814</b>

**Table 7**

<b>PATHOLOGICAL OUTCOMES</b>	<b>DESCRIPTION</b>	<b>N</b>
<b>MCM</b>	<b>all MCMs</b>	<b>702</b>
	<b>Chromosomal</b>	
CHR	Chromosomal	22
CHR	Down's syndrome	44
CHR	Edward syndrome/trisomy 18	9
CHR	Klinefelter's syndrome	2
CHR	Patau syndrome/trisomy 13	6
CHR	Turner's syndrome	4
CHR	Wolff-Hirschorn syndrome	1
<b>CHR</b>	<b>all CHR</b>	<b>88</b>
	<b>Syndromes or genetic conditions</b>	
Syndrome	Marfan's syndrome	3
Syndrome	Incontinentia pigmenti	2
Syndrome	Noonan's syndrome	3
Syndrome	Goldenhar syndrome (Oculo-auriculo-vertebral syndrome)	1
Syndrome	Di George's syndrome	1
Syndrome	Tuberous sclerosis	6
Syndrome	Craniosynostosis, inherited	1
Syndrome	Congenital cataract, inherited	1
Syndrome	Congenital glaucoma, inherited	1
Syndrome	X-linked Ichthyosis	1
Syndrome	X-linked Lissencephaly	1
Syndrome	Hearing loss, bilateral, inherited	1
Syndrome	Skeletal dysplasia (achondroplastic Dwarfism)	1
Syndrome	Freeman Sheldon Syndrome (distal arthrogyriposis type 2A)	1
<b>Syndromes</b>	<b>all Syndromes</b>	<b>24</b>
<b>Total</b>	<b>all cases with pathological outcomes</b>	<b>814</b>

In 519 out of 11,908 pregnancies with AED monotherapy, one or more MCMs were observed (4.4%) as opposed to 177 out of 2,799 pregnancies with AED polytherapy (6.3%), as shown in Table 8.

**Table 8. Pathological outcomes by AED treatment categories.**

(In this table, 886 spontaneous abortions have been excluded from the denominator).

	No AED	%	Monotherapy	%	Polytherapy	%	Total
<b>MCM</b>	<b>6</b>	3.6	<b>519</b>	4.4	<b>177</b>	6.3	<b>702 (4.7%)</b>
<b>CHR</b>	<b>1</b>	0.6	<b>71</b>	0.6	<b>16</b>	0.6	<b>88 (0.6%)</b>
<b>Syndromes</b>	<b>0</b>	0.0	<b>19</b>	0.1	<b>5</b>	0.2	<b>24 (0.2%)</b>
<b>No malformation</b>	<b>161</b>	95.8	<b>11,299</b>	94.9	<b>2,601</b>	92.9	<b>14,061 (94.5%)</b>
<b>Total</b>	<b>168</b>	100	<b>11,908</b>	100	<b>2,799</b>	100	<b>14,875 (100%)</b>

## PUBLICATIONS

Changes in AED prescribing patterns and in rates of MCM over time in the EURAP cohort were published in *Neurology*. 2019 Aug 27;93(9):e831-e840.

Outcome regarding the eight most common monotherapies has been published in *Lancet Neurology*, April 18, 2018.

The dose-dependent risk of MCM with exposure to valproate in mono- and polytherapy has also been analysed and reported (*Neurology*, Sept 8, 2015) and so has the risk of intrauterine death in association with different treatments (*Neurology* Aug 18, 2015).

A manuscript on seizure control in pregnancies with withdrawal of or switch from valproate during 1<sup>st</sup> trimester as compared with maintained valproate treatment has been published in *Epilepsia* (*Epilepsia* 2016; 57: e173-7).

Outcome in relation to exposure to individual drugs or specific drug combinations is not included in the present report.

## ORGANISATION, FUNDING AND SUPPORT

EURAP is a consortium of independent research groups working on a non-profit basis. The project is administratively organised by the Central Project Commission (CPC) with members representing different geographical areas and disciplines. The project has been supported over the years by educational grants/donations to EURAP from Bial, Eisai Pharmaceuticals, GlaxoSmithKline, GW Pharmaceuticals Janssen-Cilag, Johnson & Johnson, Novartis, Pfizer, Sanofi, Teva and UCB biopharma. In addition, national and regional networks may receive support from the same or other pharmaceutical companies.

# APPENDIX

## **Central Project Commission**

Dina Battino, Milano

Erminio Bonizzoni, Pavia

John Craig, Belfast

Dick Lindhout, Utrecht

Emilio Perucca, Pavia

Anne Sabers, Copenhagen

Sanjeev V Thomas, Trivandrum

Torbjörn Tomson, Stockholm, (chair)

Frank Vajda, Melbourne

## **Central Study Coordinator**

Dina Battino, Milan

## **Scientific Advisory Board**

Bernd Schmidt, Freiburg

Martin J Brodie, Glasgow

## **Outcome Assessment Committee**

*(The persons below have contributed to the work of the OAC during different time periods of the project)*

Chiara Pantaleoni, Milan, Italy

Claudia Ciaccio, Milan, Italy

Elisabeth Robert-Gnansia, Lyon, France

Francesca Faravelli, Genoa, Italy

Richard Finnell, Houston, Texas